

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 125 947 A1

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:

22.08.2001 Bulletin 2001/34

(51) Int Cl.⁷: **C07K 14/47, C07K 16/18,**

C12N 15/12, A61K 38/17,
A61K 39/395

(21) Application number: **99951188.4**

(22) Date of filing: **01.11.1999**

(86) International application number:

PCT/JP99/06076

(87) International publication number:

WO 00/26250 (11.05.2000 Gazette 2000/19)

(84) Designated Contracting States:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

(30) Priority: **30.10.1998 JP 31140898**

(71) Applicants:

- **Japan Science and Technology Corporation**
Kawaguchi-shi, Saitama 332-0012 (JP)
- **Arai, Kenichi**
Meguro-ku, Tokyo 153-0063 (JP)
- **Masai, Hisao**
Minatu-ku, Tokyo 108-0073 (JP)

(72) Inventors:

- **ARAI, Kenichi**
Tokyo 153-0063 (JP)
- **MASAI, Hisao**
Minatu-ku Tokyo 108-0073 (JP)

(74) Representative: **Webber, Philip Michael**

Frank B. Dehn & Co.,
179 Queen Victoria Street
London EC4V 4EL (GB)

(54) **HUMAN H37 PROTEINS AND cDNAS ENCODING THE SAME**

(57) This application provides human H37 protein having an amino acid sequence of SEQ ID NO: 1 or NO: 2; human gene encoding the protein; cDNA of the human gene which has a base sequence of SEQ ID NO;

3 or NO: 4; DNA fragment comprising a partial sequence of the cDNA; recombinant vector having the cDNA; antibody against human H37 protein; and a method for controlling the proliferation of cells by introducing the above DNA or antibody into the cells.

EP 1 125 947 A1

Description**Technical Field**

[0001] This application relates to human H37 protein and to cDNA encoding the protein. More particularly, this application relates to human H37 protein that is an activity-controlling subunit for the protein Cdc7 controlling the replication of human cells; to human gene encoding the protein; to an antibody to the H37 protein; and to a method for controlling the proliferation of human cells using such genetic engineering material and antibody.

Background Art

[0002] Proliferation of cells is initiated when a liquid factor called a growth factor is bound to the receptor on the cell surface and a signal for proliferation is transmitted into the cell. Accordingly, for an artificial induction of proliferation of incubated cells, methods where an excessive amount of a growth factor is added to a cell medium, where a receptor which is not inherently owned by the said cell is expressed on the cell surface and a factor which is specific for the receptor is added to a medium, etc. have been carried out. Further, in suppressing the cell proliferation, methods where competing molecule, antagonist or the like to the receptor protein is added to the medium to suppress the binding of the receptor to the growth factor, etc. have been carried out.

[0003] On the other hand, in the case of the cell where a proliferation signal is issued by binding of the receptor to the growth factor, a cycle in which its genomic DNA is replicated, uniformly distributed to daughter cells and then divided is repeated. Such a cycle is called "cell cycle" especially for eukaryotes. The cell cycle is basically classified into four phases. Thus, S phase when chromosomal DNA is replicated; M phase when the replicated chromosome is divided by a spindle body and then cytoplasm is divided; G1 phase which is a period from M phase finishes until S phase begins; and G2 phase which is a period from S phase finishes until M phase starts. Transition from G1 phase to S phase is particularly strictly controlled and DNA replication takes place only once in S phase.

[0004] It has been confirmed from the studies in yeast and higher eukaryotic cells that cyclin-dependent kinase plays a critical role in the cell cycle progression (Nature 292:558-560, 1981; Cell 66:731-742, 1991; Nature 349:338-393, 1991; Science 257:1958-1961, 1992; Bioassays 17:471, 1995). Further, from a genetic analysis in yeast, it has been clarified that another serine/threonine kinase plays an essential role in the initial stage of S phase (G1/S transition). Thus, characterization of cdc7 mutation which was isolated as one of the cell division cycle mutants (J. Mol. Biol. 59: 183-194, 1971), it revealed that the Cdc7 protein kinase functions immediately prior to chromosomal replication and that, during S phase, that is necessary for activation of origins (Mol. Cell Biol. 6:1590-1598, 1986; Genes Dev. 15: 480-490, 1998; Genes Dev. 15:491-501, 1998). It has been also clarified that the Cdc7 kinase activity is dependent upon the presence of a regulatory subunit, Dbf4 (Genetics 131:21-29, 1992; Mol. Cell. Biol. 13:2899-2908, 1993). Expression of Dbf4 is periodic and is regulated by both at the transcriptional and post-translational levels (Exp. Cell Res. 180:419-428, 1989). The increase in Cdc7 kinase activity at the G1/S boundary is at least accounted for by the elevated expression of Dbf4 in late G1 phase (Mol. Cell. Biol. 13:2899-2908, 1994; Exp. Cell Res. 180:419-428, 1989). In addition, since Dbf4 interacts with replication origins in vivo (Science 265:1243-1246, 1994), it has been suggested that the Cdc7 may trigger S phase by directly activating the replication initiation complex assembled at the origins.

[0005] In addition, the inventors of this application had already isolated kinases related to yeast Cdc7 from *Schizosaccharomyces pombe*, *Xenopus*, mouse and human (J. Biol. Chem. 273:23248-23257, 1998; EMBO J. 16:4340-4351, 1997; EMBO J. 14:3094-3104, 1995), and pointed out that eukaryotic chromosomal replication is regulated by a conserved mechanism involving this family of kinase.

[0006] From the findings in yeast and higher eukaryotes as mentioned above, it is expected that an artificial control of the cell proliferation is possible by means of the regulation of the Cdc7 kinase activity in cells, which is entirely other means than conventional methods using operation of growth factor/receptor binding.

[0007] However, the inventors had also found that the putative human homolog of Cdc7, huCdc7, possesses only a very low level of kinase activity when overexpressed in mammalian cells while a baculovirus expressed form of huCdc7 is inactive.

[0008] Under such circumstances, the inventors of this application have investigated a human cDNA library with a presumption of the presence of a regulatory subunit for human Cdc7, succeeded in isolating the cDNA encoding a novel protein which regulates the kinase activity by binding to huCdc7 and named the said protein encoded in this cDNA as H37 protein.

[0009] An object of the invention is to provide the novel protein obtained by the inventors in an industrially applicable form.

[0010] Another object of the invention is to provide a human gene encoding the protein and to provide a material for genetic operation including cDNA derived from the gene, an antibody to the protein.

[0011] Still another object of the invention is to provide a method for the artificial control of proliferation of human

cells using the above material for genetic operation.

Disclosure of the Invention

[0012] This application provides human H37 protein having an amino acid sequence of SEQ ID NO: 1 or NO: 2.

[0013] This application also provides human H37 protein having an amino acid sequence where one or more amino acid residue(s) in the amino acid sequence of SEQ ID NO: 1 or NO: 2 is/are deleted therefrom, substituted therefor or added thereto.

[0014] This application further provides a human gene encoding the above-mentioned human H37 protein; cDNA of said human gene which has a base sequence of SEQ ID NO: 3 or NO: 4; and DNA fragment comprising a partial sequence of those cDNAs.

[0015] This application furthermore provides a recombinant vector having the above-mentioned cDNA and an antibody against the human H37 protein.

[0016] This application still further provides a method for promoting the proliferation of cells, which comprises introducing the above-mentioned cDNA or the DNA fragment together with expression regulatory sequence into cell, and a method for suppressing the proliferation of cells, which comprises introducing the above-mentioned antibody into the cell.

Brief Description of Drawings

[0017] Fig. 1 is the result of a western blotting measuring the coimmunoprecipitation of H37 with huCdc7 expressed in mammalian cells. Lanes 1-4, immunoprecipitates; lanes 5-7, whole cell extract. Upper and middle panels, immunoprecipitated with anti-huCdc7 antibody No. 1; lower panel, immunoprecipitated with anti-myc antibody. Extracts were prepared from Cos7 cells transfected with huCdc7 in combination with H1 (lanes 2 and 5), H18 (lanes 3 and 6), H37 (lanes 4 and 7) or huCdc7 alone (lane 1). Western blotting was conducted with anti-myc antibody (upper panel) or an anti-huCdc7 antibody No. 1 (middle and lower panel).

[0018] Fig. 2 is a result of the western blotting measuring antibodies against H37 protein and association of huCdc7 and H37 *in vivo*. Nuclear extracts prepared from Cos7 cells transfected with myc-tagged H37 cDNA were blotted with anti-H37C (lane 1), anti-H37N (lane 2), anti-H37Cpep (lane 3) or anti-myc (lane 4) antibody. The arrow indicates the myc-tagged H37 protein, which carries 63 amino acids derived from 5' non-coding region in addition to the myc-tag. Immunoprecipitations from CEM extracts using either anti-huCdc7Cpep (lanes 5 and 6) or anti-H37Cpep (lanes 7 and 8) were separated on gel electrophoresis and blotted with huCdc7 monoclonal antibody (4A8). The symbols - and + indicates absence and presence, respectively, of each antigen peptide during immunoprecipitation. Lanes 9-13; immunoprecipitates, prepared from nuclear extracts of HeLa cells by anti-huCdc7 No.1 (lane 9), anti-huCdc7 monoclonal antibody 4A8 (lane 10), anti-H37C (lane 11), anti-H37N (lane 12) or anti-H37Cpep (lane 13), were blotted with anti-H37Cpep.

[0019] Fig. 3 shows the result of immunoprecipitation by the use of anti-huCdc7 antibody No.1 (lanes 1-5) or anti-myc antibody (lanes 6-10) from the extract of Cos7 cells transfected with myc-tagged H37 alone (lanes 1 and 6), or together with wild-type huCdc7 (lanes 2 and 7), or kinase negative huCdc7 (lanes 3 and 8). Wild-type huCdc7 alone (lanes 4 and 9) and kinase negative huCdc7 alone (lanes 5 and 10) were included as control.

[0020] Fig. 4 shows mobility shift of H37 induced by coexpression of wild-type huCdc7. Extracts were prepared from Cos7 cells expressing either wild-type or kinase negative huCdc7 together with myc-tagged H37. Immunoprecipitates with anti-huCdc7 antibody No. 1 (lanes 1 and 2) or anti-myc antibody (lanes 3 and 4) were blotted with anti-myc antibody (upper) or anti-huCdc7 antibody (lower). In all cases, samples were run on 8% SDS-PAGE.

[0021] Fig. 5 shows an amino acid sequence of the full-length H37 protein which is as same as the SEQ ID NO: 1.

[0022] Fig. 6 shows schematic representation of conserved regions between Dbf4 and H37. The double-arrowed region on Dbf4 was reported to be sufficient for interaction with huCdc7. Solid and gray double-arrowed region on H37 indicates the portion essential for interaction with huCdc7 or that sufficient for activation of huCdc7 kinase activity, respectively.

[0023] Fig. 7 is a schematic representation of N-terminal and C-terminal deletion derivatives of H37 proteins. The number at the end of each bar indicates the portion of the amino acid (corresponding to SEQ ID NO: 1) at the deletion endpoint. The striped region indicates Dbf4 motif-C.

[0024] Fig. 8 shows lacZ activity of H37 deletion derivatives in two-hybrid assay with huCdc7.

[0025] Fig. 9 shows the result of antibody coprecipitation method in which H37 deletion derivatives and huCdc7 were co-expressed in COS7 cells and a complex formation was measured.

[0026] Fig. 10 (a) is the result of northern analysis of H37 mRNA expression in various tissues, and (b) is the result of a northern analysis of H37 mRNA expression in various cancer cell lines.

[0027] Fig. 11 shows the results where WI38 cells in a resting phase are stimulated by addition of 10% of serum and

DNA contents at various stages were analyzed by means of FACS.

[0028] Fig. 12 shows the result of northern analysis of H37 and huCdc6 expression by using RNA extracted from cells of Fig. 11 (upper), and the graphs showing the relative expression amount of each mRNA (middle and lower).

[0029] Fig. 13 shows graphs showing each cell cycle fraction of human CEM cells fractionated by an elutriation method.

[0030] Fig. 14 shows the result of northern analysis for expression of H37 and cyclin E in each fraction of Fig. 13 (upper), and also shows the graphs showing the relative expression amounts of each mRNA (middle and lower).

[0031] Fig. 15 shows the result of FACS analysis for DNA content in which HeLa cells were stopped at latter G2 phase using nocodazole and then the cell cycle was synchronously shifted.

[0032] Fig. 16 shows the result northern analysis for expression of H37 and cyclin E in each cell cycle of Fig. 14 (upper), and also shows the graphs showing the relative expression amounts of each mRNA (middle and lower).

[0033] Fig. 17 shows H37 localization in cells measured by indirect immunofluorescent technique. The antibodies used are anti-H37C (A), anti-H37N (C) and a control antibody (E) while B, D and F were pictures stained with DAPI.

[0034] Fig. 18 shows kinetics of induction of DNA replication after serum stimulation of KD cells as measured by BrdU incorporation.

[0035] Fig. 19 shows the results in which KD cells, synchronized by serum starvation, were microinjected with various antibodies at 12 hours after serum stimulation, and then the rate of the cells incorporating the BrdU after 16 hours was measured. The numbers show the rate of the cells which are conducting the DNA synthesis.

[0036] Fig. 20 is microscopic pictures of examples of cells into which a mixture of anti-H37Cpep antibody and antigen peptide was microinjected. The pictures show the incorporated BrdU (upper), injected antibody (middle) and cells (lower).

Best Mode for Carrying Out the Invention

[0037] The human H37 protein of this invention has the amino acid sequence of SEQ ID NO: 1 and is a protein molecule encoded in the sequence region from 518th to 2541st of cDNA whose base sequence is shown in SEQ ID NO: 3. The H37 protein of this invention also has the amino acid sequence of SEQ ID NO: 2 and is a protein encoded in the sequence region from 518th to 1222nd of cDNA whose base sequence is shown in SEQ ID NO: 4. SEQ ID NO: 3 and NO: 4 are cDNAs derived from mRNA transcribed from the same genomic gene but the cDNA of SEQ ID NO: 4 is in a different splicing form from SEQ ID NO: 3, and there is a deletion from 1199th to 1259th of SEQ ID NO: 3.

[0038] Those H37 proteins can be prepared by known methods such as isolation from human organs and cell lines, chemical synthesis of peptide based upon the amino acid sequence provided by this invention, and a recombinant DNA technique using cDNA fragments provided by this invention. For example, in case the H37 protein is prepared by means of a recombinant DNA technique, RNA is prepared from a vector having the cDNA fragment of this invention by an in vitro transcription and then an in vitro translation is carried out using the above as a template whereby an expression in vitro is possible. In addition, when the translational region is recombined to a suitable vector by a known method, it is possible to express the H37 protein encoded in cDNA in a large quantity in *Escherichia coli*, *Bacillus subtilis*, yeast, insect cells, animal cells, etc.

[0039] When the human H37 protein of this invention is expressed in a microorganism such as *E. coli*, the translational region of cDNA of this invention is inserted into an expression vector having, for example, promoter, ribosome binding site, cDNA cloning site, terminator, origin replicable in microorganism, etc. whereupon a recombinant expression vector is prepared, then host cell is transformed using the said expression vector and the resulting transformant is incubated whereupon H37 protein encoded in cDNA can be produced in a microorganism in a large quantity. Alternatively, it can be expressed as a fusion protein with other protein. The resulting fusion protein is cleaved by an appropriate protease whereupon only the protein portion encoded in cDNA can be obtained.

[0040] When the human H37 protein of this invention is expressed in animal cells, the translational region of cDNA of this invention is recombined with expression vector for animal cells having, for example, promoter, poly(A) addition site, and splicing site, and then introduced into the animal cells whereupon the H37 protein of this invention can be expressed in animal cells.

[0041] The human H37 protein obtained by the method as mentioned above can be used as an antigen for the preparation of antibody for suppressing the proliferation of cells through inhibition of the kinase activity of huCdc7.

[0042] In addition, as will be confirmed by the Examples later, the human H37 protein of this invention has almost no similarity in terms of structure to cyclins which have been elucidated up to now, but it may be regarded as a cyclin-like associating factor for huCdc7 kinase in such respects that its expression is regulated by a cell cycle and that, as a result of binding to an huCdc7 catalytic subunit, its kinase activity can be activated. Accordingly, since the H37 protein is believed to be a very important target factor in a signal transduction pathway for the cell proliferation induced by a growth factor, elucidation of how the expression of the H37 protein or activity thereof is regulated by the signal of cell cycle in G1/S phases is expected to provide a big and novel finding for clarifying the molecular mechanism of cell cycle

control of cell replication in animal cells.

[0043] The human H37 protein of the invention contains peptide fragments (five or more amino acid residues) containing any partial amino acid sequence of SEQ ID NO: 1 or NO: 2. Such peptide fragments can be also used as an antigen for the preparation of antibody.

[0044] The gene of the invention is a human gene encoding the above-mentioned human H37 protein and can be isolated from known genomic libraries using, for example, the cDNA of this invention or a partial sequence thereof as a probe.

[0045] The cDNA of the invention is characterized in having a base sequence of SEQ ID NO: 3 or NO: 4, and can be cloned, for example, from cDNA libraries derived from human cells. cDNA is synthesized using poly(A)⁺ RNA extracted from human cells as a template. The human cells may be either those excised from human body by a surgical operation or cell lines. The cDNA can be prepared by known methods such as Okayama-Berg method (Okayama, H and Berg, P., Mol. Cell. Biol., 2:161-170, 1982), Gubler-Hoffmann method (Gubler, U. and Hoffman, J. Gene, 25: 263-269, 1983) and Capping method (Kato, S. et al., Gene, 150:243-250, 1994).

[0046] The human H37 protein of the invention is expressed in any tissue except brain and kidney and, therefore, when the human cDNA library prepared from human cells is screened using an oligonucleotide probe synthesized based on the base sequence of cDNA in SEQ ID NO: 3 or NO: 4, it is possible to easily prepare the same clone of the invention. It is also possible that the desired cDNA is synthesized by a polymerase chain reaction (PCR) using such an oligonucleotide as a primer.

[0047] In general, polymorphism due to the difference among individuals is frequently noted in human gene. Accordingly, cDNA of SEQ ID NO: 3 or NO: 4 where one or more nucleotide(s) is/are added thereto and/or deleted therefrom and/or other nucleotide(s) is/are substituted therefor is also covered by the invention.

[0048] Similarly, the protein where one or more nucleotide(s) is/are added thereto and/or deleted therefrom and/or other nucleotide(s) is/are substituted therefor as a result of those modifications is also covered by the invention so far as it has an activity of the protein having an amino acid sequence of SEQ ID NO: 1 or NO: 2. Further, variant protein where one or more nucleotide(s) is/are added thereto and/or deleted therefrom and/or other nucleotide(s) is/are substituted by an artificial means is covered by this invention as well.

[0049] The DNA fragment of the invention covers a cDNA fragment (10 bp or more) containing any partial base sequence of SEQ ID NO: 3 or NO: 4, or a cDNA fragment comprising an antisense strand thereof.

[0050] Antibody against human H37 protein of the invention can be prepared as a polyclonal antibody or a monoclonal antibody by a known method using protein per se or a partial peptide thereof.

[0051] A method for promoting the cell proliferation according to the invention is carried out by such a manner that a recombinant DNA comprising the cDNA having a base sequence of SEQ ID NO: 3 or NO: 4, or comprising a partial sequence thereof (such as a DNA fragment encoding 419 amino acid sequence region at the C terminal as shown in Example 3) and an expression-regulatory sequence (promoter and/or enhancer sequence for animal cells) is introduced into animal cells whereby the H37 protein having an amino sequence of SEQ ID NO: 1 or NO: 2 is excessively expressed in cell nuclei. Introduction of the recombinant DNA into cells can be carried out by a known method such as a calcium phosphate method, a method where ribosome and erythrocyte ghost are used, an electroporation method, a method where retrovirus or adenovirus is used as a vector and a microinjection method using a glass pipette. Promotion of cell proliferation as such is useful, for example, in a large-scale preparation of the stem cell that is useful for the therapy of human diseases. Stem cells can be differentiated into other kinds of cells such as blood stem cell and nerve stem cell, and are able to produce a large number of cells of different cell type that constitute the human body. Therefore, transplantation of stem cell in the diseases such as leukemia is a very important therapeutic means. However, since no liquid factor for self-proliferation of human stem cells without differentiation has been identified yet, the preparation of stem cells in an amount sufficient for the therapy has not been easy. According to the method of this invention, it could be possible that, as a result of manipulating the proliferation program in stem cells, the stem cells can be unlimitedly self-replicated and self-proliferated in vitro. In addition, such a promotion of cell proliferation in vitro could be also useful for the preparation of a large quantity of cells for introduction of gene for genetic therapy by ex vivo means.

[0052] The method for the suppression of cell proliferation according to the invention is carried out by injecting the above-mentioned antibody into cells. Alternatively, it may be carried out by inhibiting the expression of the cell-intrinsic H37 protein gene as well. An example thereof is a method where DNA encoding the ribozyme sequence or the antisense sequence to the transcript of the gene is introduced into the cells. Suppression of the cell proliferation as such is expected to provide a novel means for suppressing an excessive proliferation of cancer cells for example.

Examples

[0053] The invention will now be further illustrated by way of the following examples although the invention is not limited to those examples.

Example 1.

[0054] H37 cDNA labeled with a Flag tag was ligated to the down stream of an SRalpha promoter and introduced into IL-3 (interleukin 3)-dependent proB cell, Ba/F3, to give a stable transformant cell line constitutively expressing the Flag-tagged H37 protein. In this cell line, rate of the cell at the late S phase or G2 phase where DNA synthesis was almost finishing or was finished in the population of the proliferating cells increased as compared with the control parental cell line. On the other hand, the parental cells stopped its proliferation by removal of IL-3 and came into G0 phase but, in the stable transformant cell line, rate of the cells existing in S phase significantly increased even after the removal of IL-3. This result shows that the constitutive production of the H37 protein promotes the shift from G0/G1 phase to S phase, or the progress of S phase itself. It also suggests that the progress of cell cycle can be operated by an artificial expression of the H37 protein or a derivative thereof.

Example 2.

[0055] cDNA library of HeLa cell was prepared using a pGAD-GH vector and each vector was introduced into yeast strain CG1945 that has a recombinant plasmid with huCdc7 fused to DNA-binding domain of Gal4. As a result of a screening of about 3×10^5 cells of transformed yeast, five clones where β -galactosidase was positive were obtained. DNA base sequence of the insert was determined and the database was searched whereupon all of them were novel cDNA. Among them, three clones were same having a base sequence of SEQ ID NO: 1. This cDNA was named H37. Other two were single clones and were named H1 and H18, respectively.

[0056] The interaction of huCdc7 with protein encoded in those positive clones was further investigated using a production increase system in animal cells. Thus, each expression plasmid of H1, H18 and H37 tagged with myc epitope was transfected to animal cell Cos7 together with a full-length huCdc7 expression plasmid. The result is as shown in Fig. 1. Although the H37 protein was coimmunoprecipitated with an antibody against huCdc7, H1 and H18 proteins were not coprecipitated (cf. lanes 2-4 of the upper panel, Fig. 1). On the contrary, huCdc7 was coimmunoprecipitated by using anti-myc antibody alone in the case of the cells where myc-tagged H37 is co-expressed (cf. lane 4 of the lower lane, Fig. 1). From those results, it has been confirmed that only H37 cDNA encodes a protein efficiently interacting with huCdc7.

[0057] Then, in order to investigate the intrinsic H37 protein, each antibody against N- or C-terminal regions of H37 (anti-H37N antibody or anti-H37C antibody) was prepared. Further, antibody against C-terminal oligopeptide of H37 (anti-H37Cpep antibody) and antibody against C-terminal oligopeptide of huCdc7 (anti-huCdc7Cpep antibody) were also prepared. After that, association of each antibody with intrinsic huCdc7 and H37 in cells was measured. The result is as shown in Fig. 2. Thus, all of the antibodies to H37 specifically react with the myc-tagged H37 protein of 90 kDa expressed in Cos7 cells (cf. lanes 1-4, Fig. 2). Complex prepared from human CEM cells was able to be coprecipitated using an affinity-purified anti-peptide antibody. It was confirmed by an immunoblotting using anti-huCdc7Cpep antibody that all of huCdc7 and H37 immunoprecipitates contained huCdc7 (lanes 5 and 7, Fig. 2). This interaction of H37 with huCdc7 completely disappeared when antibody and peptide used as antigen for preparing the antibody were subjected to a pre-incubation in advance (cf. lane 8, Fig. 2). In extracts of HeLa cells, both anti-Cdc7 antibody and anti-H37 antibody were able to coprecipitate one polypeptide of 80 kDa that specifically reacted with anti-H37 antibody (lanes 9-13, Fig. 2).

[0058] From the above result, it was found that intrinsic huCdc7 and H37 protein were present in cells as a complex.

Example 3.

[0059] In order to investigate whether H37 protein has an ability of activating huCdc7, huCdc7/H37 complex prepared by expression of myc-tagged H37 and wild-type or kinase negative huCdc7 in Cos cells was immunoprecipitated by anti-huCdc7 antibody or anti-myc antibody and then a kinase reaction in vitro was measured using GST-MCM3 fusion protein as a substrate. The result is as shown in Fig. 3 and in Fig. 4. In the presence of wild-type huCdc7, phosphorylation in a good efficiency was observed for the MCM3 protein in both immunoprecipitate of anti-huCdc7 antibody and immunoprecipitate of anti-myc antibody (cf. lanes 2 and 7, Fig. 3). Further, two more phosphorylated proteins were observed and they were identified as transfected huCdc7 and myc-tagged H37 (data not shown). Since the phosphorylation as such was not detected at all in kinase negative huCdc7, it was confirmed that kinase activity of huCdc7 acted in the phosphorylation. However, kinase negative huCdc7 was also able to form a complex with H37 protein (cf. lanes 3 and 8, Fig. 3; lanes 2 and 4, Fig. 4). In addition, the mobility of H37 protein on gel electrophoresis was slow when a wild-type huCdc7 was co-expressed and detected as plural bands while such a shift was not observed in kinase negative huCdc7 (cf. lanes 1 and 3, Fig. 4). The band where the mobility was slow disappeared by a treatment with phosphatase and, therefore, it was confirmed to be a perphosphorylated H37 protein (data not shown). In addition, when huCdc7 and H37 protein were co-expressed in insect cells, it was possible to reconstitute a very strong kinase

activity which was able to efficiently phosphorylate MCM2 and MCM3 proteins (data now shown).

[0060] The above result shows that H37 protein activates huCdc7 kinase and further that H37 protein itself is phosphorylated by huCdc7.

[0061] Further, under those experimental conditions, intrinsic H37 protein level was too low and in the case where only huCdc7 catalytic subunit was expressed and, therefore, the kinase activity was little (cf. lanes 4 and 9, Fig. 3). From those facts, it was confirmed that H37 protein encoded regulatory cell unit of huCdc7 and specifically activated its kinase activity.

Example 4.

[0062] An amino acid sequence of the H37 protein (SEQ ID NO: 1) was analyzed. The result was that, as shown in Fig. 5 and Fig. 6, an amino acid sequence region having a homology of 33% with budding yeast Dbf4 was found. This conserved domain (H37 motif C) was also present in the gene similar to H37 identified in mouse, fruit fly and fission yeast (cf. Fig. 6). Further, another amino acid sequence of H37 (H37 motif N) was conserved in the H37-related gene in mouse, fruit fly and fission yeast. However, the H37 motif N was conserved in a somewhat modified form on the Dbf4 protein of a budding yeast (cf. Fig. 6; data partially not shown).

[0063] In order to determine the region on H37 protein essential for binding with huCdc7, a series of deletion derivatives of H37 at N- and C-terminals as shown in Fig. 7 was prepared, each of them was expressed in yeast as a fusion protein with Gal4 activated domain and an interaction of each deletion derivative with huCdc7 was investigated by means of a two-hybrid assay. The result is as shown in Fig. 8. As a result of deletion at N-terminal, even a deletion of 255 amino acids at N-terminal did not affect the interaction with huCdc7 ($\Delta N2$). However, when 50 amino acids at N-terminal were further deleted whereupon the H37 motif C was deleted, an interaction with huCdc7 was completely lost ($\Delta N3$).

[0064] On the other hand, with regard to deletion from C-terminal, when only 20 amino acids were deleted, the binding ability with huCdc7 decreased to an extent of about 60% (ΔC). In addition, when 243 or 369 amino acids were deleted from the C-terminal ($\Delta P2$ and ΔB), the interaction lowered to an extent of about 10% of the full-length clone. $\Delta P1$ containing only 235 amino acids of N-terminal did not interact with huCdc7. However, 50 amino acids that were commonly present in AB and $\Delta N2$ were not sufficient for an efficient interaction with huCdc7 (data not shown).

[0065] A part of the above-mentioned H37 deletion derivatives was co-expressed in COS7 cells together with huCdc7 and it was confirmed by immunoprecipitation method whether it formed a complex with huCdc7 protein. The result was that, like in the result of a two-hybrid assay, it was confirmed that only H37 deletion derivatives of deltaB and deltaN2 formed a complex with huCdc7 (cf. Fig. 9).

[0066] The above result shows that H37 motif-C is essential for an interaction of H37 protein with huCdc7 catalytic subunit but shows that H37 motif C alone is insufficient. In the budding yeast, it was reported already that the region containing H37 motif C was sufficient for binding with Cdc7 (Mol. Cell. Biol. 15:6775-6782, 1996). When an in vitro kinase reaction was carried out using those deletion derivatives, it was found that, although the extent was weak, only 419 amino acids at the C-terminal containing Dbf4 motif-C was sufficient for activation of phosphorylating ability of huCdc7 (data not shown).

Example 5.

[0067] Expression patterns of H37 mRNA in various human tissues and cancer cells were investigated by northern blotting. The result is as shown in Fig. 10 (a) and (b). By an H37 cDNA-specific probe, a transcribed product of 2.5 kb was detected in all tissues except brain and kidney and also in all cancer cells. That was contrary to the fact that, in brain and kidney, huCdc7 mRNA was relatively highly expressed (EMBO J. 16:4340-4351, 1997). Among the tissues tested, expression of H37 mRNA was highest in testicles and then in thymus, and it was reported by the inventors already that both of them were the tissues where the expression of huCdc7 catalytic subunit was particularly high as well (EMBO J. 16:4340-4351, 1997). In the testicles, two different RMA bands of 6 kb and 4 kb were also detected (cf. Fig. 10 (a)) although it was ambiguous what the real substances therefor were. It was also confirmed that H37 mRNA was expressed in a very high level in nearly all cancer cells except lung cancer cell A549 (cf. Fig. 10 (b)). That shows an important role of H37 protein in the cells having an active proliferating ability.

Example 6.

[0068] In order to investigate whether expression of H37 was regulated by cell cycle, human normal fibroblast cell WI38 was synchronized with G0 phase by means of serum starvation (cf. Fig. 11), total RNA was prepared at various periods after addition of serum and the level of H37 mRNA was investigated by northern blotting. The result is as shown in Fig. 12. H37 mRNA level was low in the cell at the resting stage and, as the cells came near the boundary of G1

and S, it gradually increased and, after 20 hours from the serum addition, it became maximum. The expression pattern shown in Fig. 12 was similar to that of transcribed product of huCdc6 that is known to be induced by growth stimulation (Mol. Cell. Biol. 15:4215-4224, 1995; Proc. Natl. Acad. Sci. USA 94:142-147, 1997).

[0069] Further, in order to investigate the changes in expression of H37 in cell cycle, human CEM cells were fractionated by means of elutriation method (cf. Fig. 13) and then a northern blotting was carried out (cf. Fig. 14). It was shown by the result thereof that H37 mRNA was low at G1 phase, increased from the late G1 phase to S phase, highly maintained during S phase and somewhat decreased in G2 phase although it was still kept high. In an experiment where HeLa cells were stopped at G2 phase by nocodazole and the cell cycle was synchronously shifted (cf. Fig. 15), it was also found that H37 mRNA decreased together with shift from G2 to G1 and increased again at shift to S phase (cf. Fig. 16). Similarly, expression of Cdc6 rose together with shift from G1 phase to S phase but there was a difference from H37 in that, as S phase proceeded, it decreased and, at G2 phase it was suppressed to low. The result shows that the expression of H37 mRNA changes even in the proliferating cell cycle, becomes maximum at S phase where it functions and is kept high throughout S phase.

[0070] After that, in order to investigate the changes in H37 protein and huCdc7 kinase activity depending upon H37 in cell cycle, amount of H37 protein in K562 cells synchronized with nocodazole was analyzed by western analysis while Cdc7 kinase activity was analyzed by phosphorylation reaction of GST-huMCM2 protein using anti-huCdc7 immunoprecipitate in the same manner as in Fig. 15. Although H37 protein was not detected from G1/M boundary to G1 phase, it started to be detected at the G1/S boundary and detected throughout S phase. This was in contrast to the fact that cyclin E protein is detected already in G1 phase and, as S phase proceeds, the amount suddenly decreases. On the other hand, huCdc kinase activity was also detected in S phase only corresponding to the level of H37 protein. Further, H37 protein was detected as plural phosphorylated bands in S phase and was thought to be phosphorylated by huCdc7 that was probably associated. On the contrary, level of huCdc7 protein was almost constant throughout cell cycle. This result proves that the level of huCdc7 protein varies in cell cycle and is kept high in S phase whereby huCdc7 kinase activity is also kept high in S phase.

[0071] In order to further investigate the expression of H37, intracellular localization of H37 protein in animal cells was measured. The result of indirect fluorescent antibody method using two kinds of H37-specific antibodies was that, in both HeLa and WI38 cells, intrinsic H37 protein was observed as very clear various spots in the nuclei (cf. Fig. 17).

[0072] When those results and the intranuclear localization of huCdc7 catalytic subunit already reported by the inventors (EMBO J. 16:4340-4351, 1997) were taken into consideration, it was confirmed that huCdc7/H37 complex was a kinase localized in the nuclei and that its regulatory subunit H37 was expressed depending upon the stage of cell cycle.

Example 7.

[0073] Function of the intrinsic H37 protein in the shift from G1 to S in cell cycle was investigated by antibody microinjection method. Antibody against N-terminal 305 amino acids of H37 protein (anti-H37N antibody) and antibody against C-terminal oligopeptide (anti-H37Cpep antibody) were treated by affinity purification and those antibodies were microinjected into normal fibroblast cells (KD cells) derived from human lip. KD cells were previously arrested at G0 phase by means of serum starvation and then synchronously proceeded in cell cycle by re-addition of serum. Numbers of the BrdU-positive cells, which BrdU is a nucleotide derivative and is incorporated into the cells, were measured whereby it was investigated how much fractions of cells at S phase were present after various stages after addition of serum. The result is as shown in Fig. 18. It was confirmed that the cells started DNA synthesis after about 18 hours from serum addition and that, after 24 hours, almost 90% of the cells were in S phase.

[0074] Accordingly, in this Example, antibody was microinjected at the period of after 12 hours after the addition of serum where the cells were still in G1 phase and, at 26 hours when cells would be of in S phase, the cells were fixed and the BrdU-positive cells were measured. The result is as shown in Fig. 19. Although 70% of the cells into which anti-H37N antibody was microinjected did not shift into S phase, there was nearly no effect by the control antibody. Even by the microinjection of the anti-H37Cpep antibody, an effect of inhibiting the shift to S phase which was same as or even more than the anti-H37N antibody was observed. In addition, when anti-H37Cpep antibody and a peptide which was an antigen for the preparation of anti-H37Cpep antibody were simultaneously microinjected into the cells, 70% or more of the cells were shifted to S phase.

[0075] Fig. 20 is a staining example of BrdU and microinjected antibody. The antibody was microinjected at the stage when the cells were from the mid to late S phase and, during that period, expression of H37 protein is believed to be low. The microinjected antibody efficiently bound to the freshly synthesized H37 protein and, as a result, it is speculated to inhibit the transfer of H37 protein into nucleus. Those results strongly suggest the function of H37, i.e. the function of huCdc7/H37 complex is required for the progression of S phase of the animal cells.

Industrial Applicability

[0076] As fully illustrated hereinabove, the present invention provided human H37 protein, a regulatory subunit for Cdc7 that regulate replication of human cells, human gene encoding this protein as well as cDNA thereof, an antibody
5 against H37 protein, and a method for controlling the proliferation of human cells using those genetic engineering materials and antibody. As a result, it provides potentially novel means for preparation of the necessary amount of stem cells or the like to be used for the therapy of various human diseases or for the suppression of proliferation of cancer cells.

10

15

20

25

30

35

40

45

50

55

SEQUENCE LISTING

<110> Japan Science and Technology Corporation

<120> Juman H37 protein and cDNA encoding the protein

<140> PCT/JP99/06067

<141> 1999-11-01

<150> JP No. 10-311408

<151> 1998-10-30

<160> 4

<210> 1

<211> 674

<212> PRT

<213> Homo sapiens

<400> 1

Met Asn Ser Gly Ala Met Arg Ile His Ser Lys Gly His Phe Gln Gly

1 5 10 15

Gly Ile Gln Val Lys Asn Glu Lys Asn Arg Pro Ser Leu Lys Ser Leu

20 25 30

Lys Thr Asp Asn Arg Pro Glu Lys Ser Lys Cys Lys Pro Leu Trp Gly

35 40 45

Lys Val Phe Tyr Leu Asp Leu Pro Ser Val Thr Ile Ser Glu Lys Leu

50 55 60

Gln Lys Asp Ile Lys Asp Leu Gly Gly Arg Val Glu Glu Phe Leu Ser

65 70 75 80

Lys Asp Ile Ser Tyr Leu Ile Ser Asn Lys Lys Glu Ala Lys Phe Ala

85 90 95

Gln Thr Leu Gly Arg Ile Ser Pro Val Pro Ser Pro Glu Ser Ala Tyr

100 105 110

	Thr	Ala	Glu	Thr	Thr	Ser	Pro	His	Pro	Ser	His	Asp	Gly	Ser	Ser	Phe	
5							115			120						125	
	Lys	Ser	Pro	Asp	Thr	Val	Cys	Leu	Ser	Arg	Gly	Lys	Leu	Leu	Val	Glu	
							130			135						140	
10	Lys	Ala	Ile	Lys	Asp	His	Asp	Phe	Ile	Pro	Ser	Asn	Ser	Ile	Leu	Ser	
							145						155			160	
	Asn	Ala	Leu	Ser	Trp	Gly	Val	Lys	Ile	Leu	His	Ile	Asp	Asp	Ile	Arg	
15								165				170				175	
	Tyr	Tyr	Ile	Glu	Gln	Lys	Lys	Lys	Glu	Leu	Tyr	Leu	Leu	Lys	Lys	Ser	
20							180				185					190	
	Ser	Thr	Ser	Val	Arg	Asp	Gly	Gly	Lys	Arg	Val	Gly	Ser	Gly	Ala	Gln	
							195			200				205			
25	Lys	Thr	Arg	Thr	Gly	Arg	Leu	Lys	Lys	Pro	Phe	Val	Lys	Val	Glu	Asp	
							210			215				220			
30	Met	Ser	Gln	Leu	Tyr	Arg	Pro	Phe	Tyr	Leu	Gln	Leu	Thr	Asn	Met	Pro	
							225					235				240	
	Phe	Ile	Asn	Tyr	Ser	Ile	Gln	Lys	Pro	Cys	Ser	Pro	Phe	Asp	Val	Asp	
35							245					250				255	
	Lys	Pro	Ser	Ser	Met	Gln	Lys	Gln	Thr	Gln	Val	Lys	Leu	Arg	Ile	Gln	
40							260				265				270		
	Thr	Asp	Gly	Asp	Lys	Tyr	Gly	Gly	Thr	Ser	Ile	Gln	Leu	Gln	Leu	Lys	
							275				280				285		
45	Glu	Lys	Lys	Lys	Lys	Gly	Tyr	Cys	Glu	Cys	Cys	Leu	Gln	Lys	Tyr	Glu	
							290				295				300		
	Asp	Leu	Glu	Thr	His	Leu	Leu	Ser	Glu	Gln	His	Arg	Asn	Phe	Ala	Gln	
50							305					315				320	
	Ser	Asn	Gln	Tyr	Gln	Val	Val	Asp	Asp	Ile	Val	Ser	Lys	Leu	Val	Phe	
55							325					330				335	

	Asp Phe Val Glu Tyr Glu Lys Asp Thr Pro Lys Lys Lys Arg Ile Lys		
	240	245	350
5	Tyr Ser Val Gly Ser Leu Ser Pro Val Ser Ala Ser Val Leu Lys Lys		
	355	360	365
10	Thr Glu Gln Lys Glu Lys Val Glu Leu Gln His Ile Ser Gln Lys Asp		
	370	375	380
15	Cys Gln Glu Asp Asp Thr Thr Val Lys Glu Gln Asn Phe Leu Tyr Lys		
	385	390	395
	Glu Thr Gln Glu Thr Glu Lys Lys Leu Leu Phe Ile Ser Glu Pro Ile		
20	405	410	415
	Pro His Pro Ser Asn Glu Leu Arg Gly Leu Asn Glu Lys Met Ser Asn		
	420	425	430
25	Lys Cys Ser Met Leu Ser Thr Ala Glu Asp Asp Ile Arg Gln Asn Phe		
	435	440	445
30	Thr Gln Leu Pro Leu His Lys Asn Lys Gln Glu Cys Ile Leu Asp Ile		
	450	455	460
	Ser Glu His Thr Leu Ser Glu Asn Asp Leu Glu Glu Leu Arg Val Asp		
35	465	470	475
	His Tyr Lys Cys Asn Ile Gln Ala Ser Val His Val Ser Asp Phe Ser		
	485	490	495
40	Thr Asp Asn Ser Gly Ser Gln Pro Lys Gln Lys Ser Asp Thr Val Leu		
	500	505	510
45	Phe Pro Ala Lys Asp Leu Lys Glu Lys Asp Leu His Ser Ile Phe Thr		
	515	520	525
	His Asp Ser Gly Leu Ile Thr Ile Asn Ser Ser Gln Glu His Leu Thr		
50	530	535	540
	Val Gln Ala Lys Ala Pro Phe His Thr Pro Pro Glu Glu Pro Asn Glu		
55	545	550	555
			560

Cys Asp Phe Lys Asn Met Asp Ser Leu Pro Ser Gly Lys Ile His Arg
 565 570 575
 5 Lys Val Lys Ile Ile Leu Gly Arg Asn Arg Lys Glu Asn Leu Glu Pro
 580 585 590
 10 Asn Ala Glu Phe Asp Lys Arg Thr Glu Phe Ile Thr Gln Glu Glu Asn
 595 600 605
 15 Arg Ile Cys Ser Ser Pro Val Gln Ser Leu Leu Asp Leu Phe Gln Thr
 610 615 620
 Ser Glu Glu Lys Ser Glu Phe Leu Gly Phe Thr Ser Tyr Thr Glu Lys
 20 625 630 635 640
 Ser Gly Ile Cys Asn Val Leu Asp Ile Trp Glu Glu Glu Asn Ser Asp
 645 650 655
 25 Asn Leu Leu Thr Ala Phe Phe Ser Ser Pro Ser Thr Ser Thr Phe Thr
 660 665 670
 30 Gly Phe
 674
 <210> 2
 35 <211> 234
 <212> PRT
 <213> Homo sapiens
 40 <400> 2
 Met Asn Ser Gly Ala Met Arg Ile His Ser Lys Gly His Phe Gln Gly
 45 1 5 10 15
 Gly Ile Gln Val Lys Asn Glu Lys Asn Arg Pro Ser Leu Lys Ser Leu
 20 25 30
 50 Lys Thr Asp Asn Arg Pro Glu Lys Ser Lys Cys Lys Pro Leu Trp Gly
 35 40 45
 55 Lys Val Phe Tyr Leu Asp Leu Pro Ser Val Thr Ile Ser Glu Lys Leu

50 55 60
 Gln Lys Asp Ile Lys Asp Leu Gly Gly Arg Val Glu Glu Phe Leu Ser
 5 65 70 75 80
 Lys Asp Ile Ser Tyr Leu Ile Ser Asn Lys Lys Glu Ala Lys Phe Ala
 10 85 90 95
 Gln Thr Leu Gly Arg Ile Ser Pro Val Pro Ser Pro Glu Ser Ala Tyr
 100 105 110
 15 Thr Ala Glu Thr Thr Ser Pro His Pro Ser His Asp Gly Ser Ser Phe
 115 120 125
 Lys Ser Pro Asp Thr Val Cys Leu Ser Arg Gly Lys Leu Leu Val Glu
 20 130 135 140
 Lys Ala Ile Lys Asp His Asp Phe Ile Pro Ser Asn Ser Ile Leu Ser
 25 145 150 155 160
 Asn Ala Leu Ser Trp Gly Val Lys Ile Leu His Ile Asp Asp Ile Arg
 165 170 175
 30 Tyr Tyr Ile Glu Gln Lys Lys Lys Glu Leu Tyr Leu Leu Lys Lys Ser
 180 185 190
 35 Ser Thr Ser Val Arg Asp Gly Gly Lys Arg Val Gly Ser Gly Ala Gln
 195 200 205
 Lys Thr Arg Thr Gly Arg Leu Lys Lys Pro Phe Val Lys Val Glu Asp
 40 210 215 220
 Met Ser Gln Ser Pro Ala Val His Leu Met
 225 230 234
 45
 <210> 3
 <211> 2780
 50
 <212> DNA
 <213> Homo sapiens
 <400> 3
 55

AATTCGGCAC GAGCTCTCTG AGGCTGCGCC AAGACCTGAA GCGGCGGACC GAGAGCCCGG 60
 GTCTGAGACT GAGAGAGCAA CGGAATGGAG GCGGGGTAGA GGCGGAAACA CAACCTGCAG 120
 5 GGCCAGAGCG AGGCGCCAGA AGGACGGCGG CGTGAGGGGG CCGGGCGCGC AGCGCGAGAA 180
 GGCAGGCACG AGGGGCGAGC GCGAGGCGGG GCACGGCGCG TGGCGTGAGA CGGGGCGGGG 240
 10 CGCGCGTATC GGCGCCCGG CCGCGTGACG CGTTTTCAA TCTTCAACCG CCGCAGCCCA 300
 CTCGTTTGTG CTTTGCGCCT TCCTCCTCCG CGCCTTGGAG CCGGATCCGG CCCCAGAAAC 360
 CCGACCTGCA GACGCGGTAC CTCTACTGCG TAGAGGCCGT AGCTGGCGGA AGGAGAGAGG 420
 15 CGGCCGTCTT GTCAACAGGC CGGGGGAAGC CGTGCTTTTC CGGCTGCCCG GTGCGACACT 480
 TTCTCCGGAC CCAGCATGTA GGTGCCGGGC GACTGCCATG AACTCCGGAG CCATGAGGAT 540
 CCACAGTAAA GGACATTTCC AGGGTGAAT CCAAGTCAA AATGAAAAA ACAGACCATC 600
 TCTGAAATCT CTGAAACTG ATAACAGGCC AGAAAAATCC AAATGTAAGC CACTTTGGGG 660
 AAAAGTATTT TACCTTGACT TACCTTCTGT CACCATATCT GAAAAACTTC AAAAGGACAT 720
 25 TAAGGATCTG GGAGGGCGAG TTGAAGAATT TCTCAGCAA GATATCAGTT ATCTTATTTT 780
 AAATAAGAAG GAAGCTAAAT TTGCACAAAC CTTGGGTGCA ATTTCTCCTG TACCAAGTCC 840
 AGAATCTGCA TATACTGCAG AAACCACTTC ACCTCATCCC AGCCATGATG GAAGTTCATT 900
 30 TAAGTCACCA GACACAGTGT GTTTAAGCAG AGGAAAATTA TTAGTTGAAA AAGCTATCAA 960
 GGACCATGAT TTTATTCCTT CAAATAGTAT ATTATCAAAT GCCTTGTCAT GGGGAGTAAA 1020
 35 AATTCTTCAT ATTGATGACA TTAGATACTA CATTGAACAA AAGAAAAAG AGTTGTATTT 1080
 ACTCAAGAAA TCAAGTACTT CAGTAAGAGA TGGGGGCAAA AGAGTTGGTA GTGGTGCACA 1140
 AAAACAAGA ACAGGAAGAC TCAAAAAGCC TTTTGTAAG GTGGAAGATA TGAGCCAACT 1200
 40 TTATAGGCCA TTTTATCTTC AGCTGACCAA TATGCCTTTT ATAAATTATT CTATTCAGAA 1260
 GCCCTGCAGT CCATTTGATG TAGACAAGCC ATCTAGTATG CAAAAGCAA CTCAGGTAA 1320
 45 ACTAAGAATC CAAACAGATG GCGATAAGTA TGGTGAACC TCAATTCAAC TCCAGTTGAA 1380
 AGAGAAGAAG AAAAAAGGAT ATTGTGAATG TTGCTTGCAG AAATATGAAG ATCTAGAAAC 1440
 TCACCTTCTA AGTGAGCAAC ACAGAACTT TGCACAGAGT AACCAGTATC AAGTTGTTGA 1500
 50 TGATATTGTA TCTAAGTTAG TTTTGTACTT TGTGGAATAT GAAAAGGACA CACCTAAAAA 1560
 GAAAAGAATA AAATACAGTG TTGGATCCCT TTCTCCTGTT TCTGCAAGTG TCCTGAAAAA 1620
 55 GACTGAACAA AAGGAAAAAG TGGAATTGCA ACATATTTCT CAGAAAGATT GCCAGGAAGA 1680

TGATACAACA GTGAAGGAGC AGAATTTCCCT GTATAAGAG ACCCAGGAAA CTGAAAAAAA 1740
 GCTCCTGTTT ATTTGAGAGC CCATCCCCCA CCCTTCAAAT GAATTGAGAG GGCTTAATGA 1800
 5 GAAAATGAGT AATAAATGTT CCATGTTAAG TACAGCTGAA GATGACATAA GACAGAATTT 1860
 TACACAGCTA CCTCTACATA AAAACAAACA GGAATGCATT CTTGACATTT CCGAACACAC 1920
 10 ATTAAGTGAA AATGACTTAG AAGAACTAAG GGTAGATCAC TATAAATGTA ACATACAGGC 1980
 ATCTGTACAT GTTTCTGATT TCAGTACAGA TAATAGTGA TCTCAACCAA AACAGAAGTC 2040
 AGATACTGTG CTTTTTCCAG CAAAGGATCT CAAGGAAAAG GACCTTCATT CAATATTTAC 2100
 15 TCATGATTCT GGTCTGATAA CAATAAACAG TTCACAAGAG CACCTAACTG TTCAGGCAAA 2160
 GGCTCCATTG CATACTCCTC CTGAGGAACC CAATGAATGT GACTTCAAGA ATATGGATAG 2220
 20 TTTACCTTCT GGTAATAAC ATCGAAAAGT GAAAATAATA TTAGGACGAA ATAGAAAAGA 2280
 AAATCTGGAA CCAAATGCTG AATTTGATAA AAGAACTGAA TTTATTACAC AAGAAGAAAA 2340
 CAGAATTTGT AGTTCACCGG TACAGTCTTT ACTAGACTTG TTTGAGACTA GTGAAGAGAA 2400
 25 ATCAGAATTT TTGGGTTTCA CAAGCTACAC AGAAAAGAGT GGTATATGCA ATGTTTTAGA 2460
 TATTTGGGAA GAGGAAAATT CAGATAATCT GTTAACAGCG TTTTCTCGT CCCCTTCAAC 2520
 TTCTACATTT ACTGGCTTTT AGAATTTAAA AAATGCATAC TTTTCAGAAG TGATAAGGAT 2580
 30 CATATTCTTG AAATTTTAT AAATATGTAT GGAAATTCCT AGGATTTTTT TACCAGCTTT 2640
 GTTTACAGAC CCAAATGTAA ATATTAATAA TAAATATTG CAATTTTCTA CAGAATTGAA 2700
 35 TACCTGTAA AGAAAAATTA CAGAATAAAC TTGTGACTCG TCTTGTTTTA CATTAAAAAA 2760
 AAAAAAAAAA AAAACTCGAG 2780

<210> 4

<211> 2719

<212> DNA

<213> Homo sapiens

<400> 4

AATTGGCAC GAGCTCTCTG AGGCTGCGCC AAGACCTGAA GCGGCGGACC GAGAGCCCGG 60
 50 GTCTGAGACT GAGAGAGCAA CGGAATGGAG GCGGGGTAGA GGCGGAAACA CAACCTGCAG 120
 GGCCAGAGCG AGGCGCGAGA AGGACGGCGG CGTCAGGGGG CGGGGCGCGC AGCGCGAGAA 180
 55 GGCAGGCACG AGGGGCGAGC GCGAGGGGGG GCACGGCGCG TGGCGTGAGA CGGGGCGGGG 240

CGCGCGTATC GGCGCGCGG CGCGGTGACG CGTTTTCAA TCTTCAACCG CCGCAGCCCA 300
 CTCGTTTGTG CTTTGC GCCT TCCTCCTCCG CGCCTTGGAG CCGGATCCGG CCCC GGAAAC 360
 CCGACCTGCA GACGCGGTAC CTCTACTGCG TAGAGGCCGT AGCTGGCGGA AGGAGAGAGG 420
 CGGCCGTCTT GTCAACAGGC CGGGGGAAGC CGTGCTTTTC CGCCTGCCCC GTGCGACACT 480
 TTCTCCGGAC CCAGCATGTA GGTGCCGGGC GACTGCCATG AACTCCGGAG CCATGAGGAT 540
 CCACAGTAAA GGACATTTCC AGGGTGAAT CCAAGTCAA AATGAAAAA ACAGACCATC 600
 TCTGAAATCT CTGAAACTG ATAACAGGCC AGAAAAATCC AAATGTAAGC CACTTTGGGG 660
 AAAAGTATTT TACCTTGACT TACCTTCTGT CACCATATCT GAAAAACTTC AAAAGGACAT 720
 TAAGGATCTG GGAGGCCGAG TTGAAGAATT TCTCAGCAA GATATCAGTT ATCTTATTTT 780
 AAATAAGAAG GAAGCTAAAT TTGCACAAAC CTTGGGTGCA ATTTCTCTG TACCAAGTCC 840
 AGAATCTGCA TATACTGCAG AAACCACTTC ACCTCATCCC AGCCATGATG GAAGTTCATT 900
 TAAGTCACCA GACACAGTGT GTTTAAGCAG AGGAAAATTA TTAGTTGAAA AAGCTATCAA 960
 GGACCATGAT TTTATTGCTT CAAATAGTAT ATTATCAAAT GCCTTGTCAT GGGGAGTAAA 1020
 AATTCTTCAT ATTGATGACA TTAGATACTA CATTGAACAA AAGAAAAAG AGTTGTATTT 1080
 ACTCAAGAAA TCAAGTACTT CAGTAAGAGA TGGGGGCAA AGAGTTGGTA GTGGTGCACA 1140
 AAAACAAGA ACAGGAAGAC TCAAAAAGCC TTTTGTAAG GTGGAAGATA TGAGCCAAAG 1200
 CCCTGCAGTC CATTTGATGT AGACAAGCCA TCTAGTATGC AAAAGCAAAC TCAGGTAAAA 1260
 CTAAGAATCC AAACAGATGG CGATAAGTAT GGTGGAACCT CAATTCAACT CCAGTTGAAA 1320
 GAGAAGAAGA AAAAAGGATA TTGTGAATGT TGCTTGCAAG AATATGAAGA TCTAGAACT 1380
 CACCTTCTAA GTGAGCAACA CAGAACTTT GCACAGAGTA ACCAGTATCA AGTTGTTGAT 1440
 GATATTGTAT CTAAGTTAGT TTTTGACTTT GTGGAATATG AAAAGGACAC ACCTAAAAAG 1500
 AAAAGAATAA AATACAGTGT TGGATCCCTT TCTCCTGTTT CTGCAAGTGT CCTGAAAAAG 1560
 ACTGAACAAA AGGAAAAAGT GGAATTGCAA CATATTTCTC AGAAAGATTG CCAGGAAGAT 1620
 GATACAACAG TGAAGGAGCA GAATTTCTG TATAAAGAGA CCCAGGAAAC TGAAAAAAG 1680
 CTCCTGTTTA TTTGAGAGC CATCCCCAC CTTCAAATG AATTGAGAGG GCTTAATGAG 1740
 AAAATGAGTA ATAAATGTTG CATGTAAAGT ACAGCTGAAG ATGACATAAG ACAGAATTTT 1800
 ACACAGCTAC CTCTACATAA AAACAAACAG GAATGCATTC TTGACATTTT CGAACACACA 1860
 TTAAGTCAAA ATGACTTAGA AGAACTAAGG GTAGATCACT ATAAATGTAA CATACAGGCA 1920

TCTGTACATG TTTCTGATTT CAGTACAGAT AATAGTGGAT CTCAACCAAA ACAGAAGTCA 1980
 GATACTGTGC TTTTCCAGC AAAGGATCTC AAGGAAAAGG ACCTTCATTC AATATTTACT 2040
 CATGATTCTG GTCTGATAAC AATAACAGT TCACAAGAGC ACCTAACTGT TCAGGCAAAG 2100
 GCTCCATTCC ATACTCCTCC TGAGGAACCC AATGAATGTG ACTTCAAGAA TATGGATAGT 2160
 TTACCTTCTG GTAAATACA TCGAAAAGTG AAAATAATAT TAGGACGAAA TAGAAAAGAA 2220
 AATCTGGAAC CAAATGCTGA ATTTGATAAA AGAACTGAAT TTATTACACA AGAAGAAAAC 2280
 AGAATTTGTA GTTCACCGGT ACAGTCTTTA CTAGACTTGT TTCAGACTAG TGAAGAGAAA 2340
 TCAGAATTTT TGGGTTTCAC AAGCTACACA GAAAAGAGTG GTATATGCAA TGTTTATAGT 2400
 ATTTGGAAG AGGAAAATTC AGATAATCTG TTAACAGCGT TTTTCTCGTC CCCTTCAACT 2460
 TCTACATTTA CTGGCTTTTA GAATTTAAAA AATGCATACT TTTCAGAAGT GATAAGGATC 2520
 ATATTCTTGA AATTTTATA AATATGTATG GAAATTCTTA GGATTTTTTT ACCAGCTTTG 2580
 TTTACAGACC CAAATGTAAA TATTAATAAT AAATATTTGC AATTTTCTAC AGAATTGAAT 2640
 ACCTGTTAAA GAAAAATTAC AGAATAAACT TGTGACTGGT CTTGTTTTAC ATTAACAAAA 2700
 AAAAAAAAAA AAACCTCGAG 2719

Claims

1. Human H37 protein having an amino acid sequence of SEQ ID NO: 1.
2. Human H37 protein having an amino acid sequence where one or more amino acid residue(s) in the amino acid sequence of SEQ ID NO: 1 is/are deleted therefrom, substituted therefor or added thereto.
3. Human H37 protein having an amino acid sequence of SEQ ID NO: 2.
4. Human H37 protein having an amino acid sequence where one or more amino acid residue(s) in the amino acid sequence of SEQ ID NO: 2 is/are deleted therefrom, substituted therefor or added thereto.
5. A human gene encoding the human H37 protein of claim 1 or 2.
6. A cDNA of the human gene of claim 5, which has a base sequence of SEQ ID NO: 3 or a base sequence where one or more base(s) in the base sequence of SEQ ID NO: 3 is/are deleted therefrom, substituted therefor or added thereto.
7. A cDNA of the human gene of claim 5, which has a base sequence of SEQ ID NO: 4 or a base sequence where one or more base(s) in the base sequence of SEQ ID NO: 4 is/are deleted therefrom, substituted therefor or added thereto.
8. A DNA fragment comprising a partial sequence of cDNA of claim 6.
9. A DNA fragment comprising a partial sequence of cDNA of claim 7.
10. A recombinant vector having the cDNA of claim 6.

11. A recombinant vector having the cDNA of claim 7.

12. An antibody against the human H37 protein of claim 1 or 2.

5 **13.** An antibody against the human H37 protein of claim 3 or 4.

14. A method for promoting the proliferation of cells, which comprises introducing the cDNA of claim 6 or the DNA fragment of claim 8 together with expression-regulatory sequences into the cells.

10 **15.** A method for promoting the proliferation of cells, which comprises introducing the cDNA of claim 7 or the DNA fragment of claim 9 together with an expression regulatory sequence into the cells.

16. A method for suppressing the proliferation of cells, which comprises introducing the antibody of claim 10 into the cells.

15 **17.** A method for suppressing the proliferation of cells, which comprises introducing the antibody of claim 11 into the cells.

20

25

30

35

40

45

50

55

Fig. 1




Fig.2







Fig.3

Fig. 4



24

Fig. 6



Dbf4 motif-N

H37 32: **LT** **T** **D** **N** **F** **E** **K** **S** **N** **G** **K** **P** **L** **H** **G** **R** **V** **E** **I** **D** **L** **P** **S** **V** **T** **S** **E** **K** **L** **O** **K** **D** **I** **K** **E** **L** **G** **G** **R** **V** **E** **E** **F** **L** **S** **K** **D** **I** **S** **Y** **L** **S** **N** **K** **K** 91
 mu-H37 : **L** **R** **A** **D** **N** **F** **E** **K** **S** **N** **G** **K** **P** **L** **H** **G** **R** **V** **E** **I** **D** **L** **P** **S** **V** **T** **S** **E** **K** **L** **O** **K** **D** **I** **K** **E** **L** **G** **G** **R** **V** **E** **E** **F** **L** **S** **K** **D** **I** **S** **Y** **F** **V** **S** **N** **K** **K**
 Dm-H37 : **T** **P** **P** **K** **V** **K** **V** **I** **K** **S** **N** **G** **K** **P** **L** **H** **G** **R** **V** **E** **I** **D** **L** **P** **S** **V** **T** **S** **E** **K** **L** **O** **K** **D** **I** **K** **E** **L** **G** **G** **R** **V** **E** **E** **F** **L** **S** **K** **D** **I** **S** **Y** **L** **S** **N** **K** **K**
 *** ** ****

huDbf4N 92: **E** **A** **R** **F** **A** **Q** **T** 98
 muDbf4N : **E** **A** **R** **F** **A** **Q** **T**
 DmDbf4N : **L** **V** **I** **G** **G** **T** **S**
 *

Dbf4 motif-C

H37 263: **K** **Q** **T** **G** **V** **K** **L** **R** **I** **Q** **T** **D** **G** **O** **K** **Y** **G** **C** **S** **I** **Q** **L** **Q** **L** **K** **E** **K** **K** **K** **G** **Y** **C** **E** **C** **C** **L** **Q** **K** **Y** **E** **D** **L** **E** **T** **H** **L** **S** **E** **C** **H** **R** **N** **F** **A** **Q** **S** **N** 322
 mu-H37 : **K** **Q** **A** **C** **P** **K** **L** **R** **I** **N** **M** **D** **G** **O** **K** **C** **G** **P** **V** **Q** **L** **Q** **L** **K** **E** **K** **R** **K** **K** **G** **Y** **C** **E** **C** **C** **L** **Q** **K** **Y** **E** **D** **L** **E** **T** **H** **L** **S** **E** **C** **H** **R** **N** **F** **A** **Q** **S** **N**
 Dm-H37 : **P** **S** **L** **Q** **E** **L** **K** **K** **Q** **S** **A** **I** **P** **N** **S** **P** **R** **S** **N** **C** **R** **E** **P** **I** **D** **S** **S** **E** **R** **Q** **C** **G** **V** **C** **E** **I** **C** **K** **L** **E** **T** **I** **L** **N** **I** **H** **I** **O** **S** **K** **D** **H** **E** **L** **F** **A** **K** **N** **S**
 Dbf4 619: **K** **K** **S** **T** **S** **T** **N** **V** **T** **L** **H** **F** **N** **A** **Q** **T** **A** **C** **A** **Q** **P** **V** **K** **K** **E** **T** **V** **K** **N** **S** **G** **Y** **C** **H** **N** **R** **V** **K** **E** **S** **L** **E** **Q** **I** **V** **S** **L** **C** **H** **L** **S** **F** **A** **E** **N** 677
 * * * * *

Fig. 7




Fig.8




Fig.9




Fig.10




Fig. 11




Fig-12





Fig.13

Fig.14






Fig.15




Fig.16






Fig. 17




Fig.18




Fig.19





Fig.20

Microinjection with



INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP99/06076

A. CLASSIFICATION OF SUBJECT MATTER Int.Cl ⁷ C07K14/47, 16/18, C12N15/12, A61K38/17, 39/395		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) Int.Cl ⁷ C07K14/00-14/825, C12N15/11-15/62		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) GenBank/EMBL/DDBJ/GeneSeq, SwissProt/PIR/GeneSeq, MEDLINE		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	Molecular and Cellular Biology, Volume 19, Number 7, issued July 1999, Hiroyuki Kumagai et al., "A Novel Growth-and Cell Cycle-Regulated Protein, ASK, Activates Human Cdc7-Related Kinase and Is Essential for G1/S Transition Mammalian Cells", pages 5083-5095	1-17
P, X	Database MEDLINE on PubMed, Accession No. 10523313, Jiang, W. et al., "Mammalian Cdc7-Dbf4 protein kinase complex is essential for initiation of DNA replication", EMBO Journal, Vol. 18, Number 20, issued 15 October 1999, pages 5703-5713	1-17
P, X	Database GenBank, Accession No. AF160876, 22 July 1999, Hollingsworth, R., "Homo sapiens DBF4-like protein (DBF4) mRNA, complete cds."	1-17
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "I" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search 25 January, 2000 (25.01.00)		Date of mailing of the international search report 08 February, 2000 (08.02.00)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP99/06076

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	The Journal of Biological Chemistry, Vol. 273, No. 36, issued 4 September 1998, Jung Min Kim et al., "Growth Regulation of the Expression of Mouse cDNA and Gene Encoding a Serine/Threonine Kinase Related to <i>Saccharomyces cerevisiae</i> CDC7 Essential for G1/S Transition", pages 23248-23257	1-17
A	Proceedings of the National Academy of Sciences, USA, Vol. 94, No. 26, issued 23 December 1997, Wei Jiang et al., "Identification and characterization of a human protein kinase related to budding yeast Cdc7p", pages 14320-14325	1-17
A	The EMBO Journal, Vol. 16, No. 14, issued 16 July 1997, Noriko Sato et al., "Human and <i>Xenopus</i> cDNAs encoding budding yeast Cdc7-related kinases: in vitro phosphorylation of MCM subunits by a putative human homologue of Cdc7", pages 4340-4351	1-17

Form PCT/ISA/210 (continuation of second sheet) (July 1992)